

**Each Random Variable in separable Banach Space
belongs to the Domain of Definition of some
inverse to compact linear non-random operator.**

E.Ostrovsky^a, L.Sirota^b

^a Corresponding Author. Department of Mathematics and computer science.
Bar-Ilan University, 84105, Ramat Gan, Israel.

E-mail: eugostrovsky@list.ru

^b Department of Mathematics and computer science.
Bar-Ilan University, 84105, Ramat Gan, Israel.

E-mail: sirota3@bezeqint.net

Abstract.

We give in this short report a very simple proof that arbitrary random variable with Borelian distribution in separable Banach space belongs with probability one to a pre-image of some linear compact non-random operator.

Key words and phrases: Law of distribution, Banach and Linear Topological Spaces, random variables (r.v.), non-random sequence, support of measure, compact imbedded subspace and compact linear operator, Franklin's functional orthonormal system and constant.

1 Introduction. Notations. Definitions. Statement of problem. Short history.

Let

A. X be arbitrary Separable Banach Space (SBS) with a norm $\|\cdot\|_X$, briefly: $X = (X, \|\cdot\|_X)$, equipped with Borelian sigma-algebra B ;

B. $(\Omega, F, \mathbf{P})$ be probability triple with expectation $\mathbf{E}$;

C. $\xi : \Omega \rightarrow X$ be fixed random variable (r.v.) defined on the our triple with the values in the space X :

$$\mathbf{P}(\xi \in X) = 1 \tag{1.1}$$

with Borelian distribution $\mu_\xi(\cdot)$:

$$\mu_\xi(A) = \mathbf{P}(\xi \in A), \quad A \in \mathcal{B}. \quad (1.2)$$

We will denote the set of all *compact* linear continuous operators acting from the Banach space X_1 with values in the Banach space X_2 as $\text{Comp}(X_1, X_2)$; we agree for brevity $\text{Comp}(X, X) := \text{Comp}(X, X)$.

The domain of definition of the inverse operator $U \in \text{Comp}(X)$ will be denoted as ordinary $\text{Dom}(U^{-1})$:

$$\text{Dom}(U^{-1}) = \{x, \ x \in X, \ \exists y \in X \Rightarrow x = Uy\}.$$

It is known, see [1], [6], [7], that there exists a subspace Y which dependent only on the distribution ξ , compact imbedded in the space X , i.e. such that the unit ball of the space Y is pre-compact set of the space X , which may be as a capacity of the support μ_ξ :

$$\mu_\xi(Y) = \mathbf{P}(\xi \in Y) = 1. \quad (1.3)$$

Our aim in this report is to generalize the equality (1.3) and consider some applications.

2 Main result.

Theorem 2.1. There exists a compact operator $U \in \text{Comp}(X)$ which dependent only on the distribution μ_ξ such that the inverse operator U^{-1} is unique defined on the set $\text{Dom}(U^{-1})$ and such that $\xi \in \text{Dom}(U^{-1})$ almost everywhere:

$$\mathbf{P}(U^{-1}(\xi) \in X) = 1. \quad (2.1)$$

Proof.

A. Let us start for beginning from the case when the space X coincides with the space c_0 of the sequences goes to zero:

$$c_0 = \{x\} = \{\vec{x}\} = \{x_1, x_2, \dots, x_n, \dots\}, \quad \lim_{n \rightarrow \infty} x_n = 0, \quad \|x\| \stackrel{\text{def}}{=} \sup_n |x_n|. \quad (2.2)$$

Let the random vector $\xi = \vec{\xi}$ belongs to the space c_0 , so that $\lim_{n \rightarrow \infty} \xi_n = 0$ a.e. There exists a strictly positive non-random sequence $\epsilon = \vec{\epsilon} = \{\epsilon_1, \epsilon_2, \dots\}$ tending to zero: $\vec{\epsilon} \in c_0$ such that

$$\xi_n = \epsilon_n \cdot \eta_n, \quad \lim_{n \rightarrow \infty} \eta_n = 0 \pmod{\mathbf{P}}, \quad (2.3)$$

see [5], chapter 7, section 4.

Define the following linear operator $U : c_0 \rightarrow c_0$ as follows:

$$U(\{x_1, x_2, \dots, x_n, \dots\}) = \{\epsilon_1 x_1, \epsilon_2 x_2, \dots, \epsilon_n x_n, \dots\}; \quad (2.4)$$

then $U \in \text{Comp}(c_0)$ and $\vec{\xi} \in \text{Dom}(U^{-1}) \pmod{\mathbf{P}}$.

B. At the same proof may be used in the case when the space X is equal to the space $c_0(Y)$, which consists by definition on all the sequences of the elements of the (non necessary to be separable) Banach space Y tending to zero:

$$c_0(Y) = \{y_i, i = 1, 2, \dots, y_i \in Y, \lim_{n \rightarrow \infty} y_n = 0\}. \quad (2.5)$$

C. The general case may be reduced to the considered in (2.5).

Note first of all that by virtue of the universality of the space $C(0, 1)$ in the class of all separable Banach spaces it is sufficient to consider only the case when the space B coincides with the classical space of all continuous functions defined on the closed interval $[0, 1]$: $B = C[0, 1]$, equipped with uniform norm $\|f\| := \max_{t \in [0, 1]} |f(t)|$, see [6].

Further, we will use the Franklin orthonormal sequence of functions $\phi_i = \phi_i(t)$, $t \in [0, 1]$, $i = 1, 2, \dots$, see [3]; the detail investigation of these functions are obtained in [4], [2]. Note only that these orthonormal functions forms the unconditional basis in the space $C(0, 1)$.

Let $\xi(t)$, $t \in [0, 1]$ be continuous with probability one random process: $\mathbf{P}(\xi(\cdot) \in C[0, 1]) = 1$. It may be expressed in the uniform convergent Fourier - Franklin series

$$\xi(t) = \sum_{i=1}^{\infty} \xi_i \phi_i(t), \quad \xi_i = \int_0^1 \xi(t) \phi_i(t) dt. \quad (2.6)$$

so that with probability one

$$\lim_{n \rightarrow \infty} \|\xi(t) - \sum_{i=1}^n \xi_i \phi_i(t)\| = 0. \quad (2.7)$$

Let $\{N(k)\}$, $k = 1, 2, \dots$ be strictly increasing integer non-random sequence, $N(1) = 1$. We can and will suppose without loss of generality that there exists a continuous strictly increasing function $N_1(x)$, defined on the semi - axis $[1, \infty)$, such that $N_1(x)/(x = k) = N(k)$. We will denote such a function as before for brevity $N(x)$.

Define a sequence of linear operators $Q_k[\cdot](t)$

$$Q_k[\xi](t) = \sum_{n=N(k)+1}^{N(k+1)} \xi_n \phi_n(t), \quad \zeta(k) = \|Q_k[\xi]\|, \quad (2.8)$$

and correspondingly for arbitrary (continuous) numerical function $f = f(t)$

$$Q_k[f](t) = \sum_{n=N(k)+1}^{N(k+1)} f_n \phi_n(t), \quad f_i = \int_0^1 f(t) \phi_i(t) dt. \quad (2.8a)$$

The sequence $\zeta = \{\zeta(k)\}$ belongs the the space c_0 almost surely.

In accordance with the item **A** of this section there exists a non-random strictly increasing unbounded sequence $w(k)$ such that by the properly of the selected sequence $\{N(k)\}$

$$\sum_k w(k)\zeta(k) < \infty \pmod{\mathbf{P}}. \quad (2.9)$$

The required operator $U : C[0, 1] \rightarrow C[0, 1]$ may be constructed as follows

$$U^{-1}[f](t) \stackrel{\text{def}}{=} \sum_{k=1}^{\infty} w(k) Q_k[f](t). \quad (2.10)$$

Indeed,

$$\begin{aligned} U^{-1}[\xi](t) &:= \sum_{k=1}^{\infty} w(k) Q_k[\xi](t) = \sum_{k=1}^{\infty} w(k) \sum_{n=N(k)+1}^{N(k+1)} \xi_n \phi_n(t) = \\ &\sum_{n=1}^{\infty} v(n) \xi_n \phi_n(t), \end{aligned}$$

where the coefficients $v(n)$ have a form

$$v(n) = \sum_{k=1}^{\infty} w(k) \cdot I(N(k) + 1 \leq n \leq N(k + 1)), \quad (2.11)$$

where $I(\cdot)$ denotes the indicator function.

Denote

$$\Delta(n) = N^{-1}(n - 1) - (N^{-1}(n) - 1). \quad (2.12)$$

Note that the values $v(n)$ allows the following bilateral estimates:

$$\begin{aligned} v_-(n) &:= w \left(\text{Ent}[N^{-1}(n) - 1] \right) \cdot (\Delta(n) - 1) \leq v(n) \leq \\ &w \left(\text{Ent}[N^{-1}(n - 1)] + 1 \right) \cdot (\Delta(n) + 1) =: v_+(n), \end{aligned} \quad (2.13)$$

where $\text{Ent}(z)$ denotes the integer part of the value z .

The operator $U[f]$ has a form

$$U[f](t) = \sum_{n=1}^{\infty} \frac{f_n \phi_n(t)}{v(n)} = \int_0^1 R(t, s) f(s) ds, \quad (2.14)$$

i.e. is linear integral compact ("diagonal") operator with continuous by means of the choosing of the sequence $\{N(k)\}$ kernel

$$R(t, s) = \sum_{n=1}^{\infty} \frac{\phi_n(t) \phi_n(s)}{v(n)}. \quad (2.15)$$

Remark 2.1. Note that the kernel, i.e. the function of two variables $R(t, s)$ in (2.15) is in addition symmetrical and positive definite; therefore the introduced operator U is self-conjugated relative the ordinary inner product

$$(g_1, g_2) := \int_0^1 g_1(t) g_2(t) dt.$$

3 Retaining of moments.

Recall that the function $\Phi : R \rightarrow R_+$ is called Young function, iff it is even, convex, continuous, non - negative, $\Phi(u) = 0 \Leftrightarrow u = 0$, is strictly increasing such that $\lim_{u \rightarrow \infty} \Phi(u) = \infty$.

By definition, $\Phi(\cdot)$ satisfies the so-called Δ_2 condition, write: $\Phi \in \Delta_2$, iff

$$\forall \lambda > 0 \Rightarrow \sup_{u > 0} \left[\frac{\Phi(\lambda u)}{\Phi(u)} \right] < \infty. \quad (3.1)$$

For instance, the functions $\Phi_p(u) = |u|^p$, $p = \text{const} > 0$ and $\Psi_{p,r}(u) = |u|^p(1 + \ln(e + |u|))^r$, $p = \text{const} > 0$, $r = \text{const}$ belong to the set Δ_2 .

The Young function $\Psi(\cdot)$ is called weaker than $\Phi(\cdot)$, if for all positive constant λ ; $\lambda = \text{const} > 0$

$$\lim_{u \rightarrow \infty} \left[\frac{\Psi(\lambda u)}{\Phi(u)} \right] = 0. \quad (3.2)$$

Notation: $\Psi \ll \Phi$.

Using for us facts about Orlicz's spaces are presented in monographs [18], [19], [20]; in this report all the Orlicz spaces are constructed over our probability space.

Theorem 3.1. Let ξ be a random vector with values in the separable Banach space $(X, \|\cdot\| = \|\cdot\|_X)$ such that for some Young function $\Phi(\cdot)$ belonging to the class Δ_2

$$\mathbf{E}\Phi(\|\xi\|_X) < \infty, \quad (3.3)$$

or equally such that the r.v. $\|\xi\|_X$ belongs to the Orlicz space $L(\Phi)$.

There exists a compact linear operator $V : X \rightarrow X$ depending only on the distribution $\text{Law}(\xi)$ and on the function $\Phi(\cdot)$ such that $\xi \in \text{Dom}(V^{-1})$ and moreover

$$\mathbf{E}\Phi(\|V^{-1}[\xi]\|_X) < \infty. \quad (3.4)$$

or equally such that the r.v. $\|V^{-1}\xi\|_X$ belongs to the Orlicz space $L(\Phi)$.

Proof. It is sufficient to consider as before the case when $X = C[0, 1]$. On the other words, let as in the second section $\xi(t)$, $t \in [0, 1]$ be continuous with probability one random process: $\mathbf{P}(\xi(\cdot) \in C[0, 1]) = 1$, and suppose in addition

$$\mathbf{E}\Phi\left(\sup_{t \in [0, 1]} |\xi(t)|\right) = \mathbf{E}\Phi(\|\xi(\cdot)\|) < \infty. \quad (3.5)$$

It may be expressed in the uniform convergent Fourier-Franklin series

$$\xi(t) = \sum_{i=1}^{\infty} \xi_i \phi_i(t), \quad \xi_i = \int_0^1 \xi(t) \phi_i(t) dt, \quad (3.6)$$

and we denote

$$F_n[f](t) = \sum_{i=1}^n f_i \phi_i(t), \quad f(\cdot) \in C[0, 1]. \quad (3.7)$$

It follows from the uniform boundedness principle that there is an absolute constant C_F (Franklin's constant) such that

$$\sup_n \|F_n[f]\| \leq C_F \|f\|, \quad (3.8)$$

so that with probability one

$$\lim_{n \rightarrow \infty} \|\xi(\cdot) - F_n[\xi](\cdot)\| = 0.$$

Further, as long as $\|\xi(\cdot) - F_n[\xi](\cdot)\| \leq (C_F + 1) \|\xi\|$, and the function $\Phi(\cdot)$ is Young function, it follows from dominated convergence theorem

$$\lim_{n \rightarrow \infty} \mathbf{E}\Phi(\|\xi - F_n[\xi]\|_c) = 0. \quad (3.9)$$

In the language of the theory of Orlicz spaces (see [18], [19], [20]) the equality (3.9) denotes the *moment convergence*, or convergence *in mean* the variable $\mathbf{E}\Phi(\|\xi - F_n[\xi]\|_c)$ to zero as $n \rightarrow \infty$.

But the function Φ satisfies the Δ_2 condition, the equality (3.9) means also the following Orlicz norm convergence:

$$\lim_{n \rightarrow \infty} \|\xi - F_n[\xi]\|_c = 0, \quad L(\Phi) = 0, \quad (3.10)$$

where for the sake of definiteness $\|f\|_c$ denotes the uniform norm $\|f\|$ and $\|\eta\|L(\Phi)$ denotes the Orlicz's norm relative the Young function Φ of the r.v. η defined on our probability space.

Let $\{M(k)\}$, $k = 1, 2, \dots$ be strictly increasing integer non - random sequence, $M(1) = 1$. We can and will suppose without loss of generality that there exists a continuous strictly increasing function $M_1(x)$, defined on the semi - axis $[1, \infty)$, such that $N_1(x)/(x = k) = M(k)$. We will denote such a function as before for brevity $M(x)$.

Define a following sequence of linear operators $R_k[\cdot](t)$

$$R_k[\xi](t) = \sum_{n=M(k)+1}^{M(k+1)} \xi_n \phi_n(t), \quad \nu(k) = \|R_k[\xi]\|_c, \quad (3.11)$$

and correspondingly for arbitrary (continuous) numerical function $f = f(t)$

$$R_k[f](t) = \sum_{n=M(k)+1}^{M(k+1)} f_n \phi_n(t), \quad f_i = \int_0^1 f(t) \phi_i(t) dt. \quad (3.12)$$

The sequence $\{M(k)\}$ may be picked such that

$$\| \nu(k) \|_{L(\Phi)} \leq 4^{-k} \| \xi(\cdot) \|_c \|_{L(\Phi)}. \quad (3.12)$$

The required operator $V : C[0, 1] \rightarrow C[0, 1]$ may be constructed as follows

$$V^{-1}[f](t) := \sum_k 2^k R_k[f](t). \quad (3.13)$$

Indeed,

$$\| \| V^{-1}[\xi] \|_c \|_{L(\Phi)} \leq \sum_k 2^{-k} \| \xi(\cdot) \|_c \|_{L(\Phi)} \leq \| \xi(\cdot) \|_c \|_{L(\Phi)}, \quad (3.14)$$

Q.E.D.

Theorem 3.2. Let ξ be again a random vector with values in the separable Banach space $(X, \| \cdot \| = \| \cdot \|_X)$ such that for some Young function $\Phi(\cdot)$ not necessary to be from the set Δ_2

$$\mathbf{E}\Phi(\| \xi \|_X) < \infty, \quad (3.15)$$

or equally such that the r.v. $\| \xi \|_X$ belongs to the Orlicz space $L(\Phi)$.

Let also $\Psi(\cdot)$ be other arbitrary Young function weaker than Φ : $\Psi \ll \Phi$. There exists a compact linear operator $V : X \rightarrow X$ depending only on the distribution $\text{Law}(\xi)$ and on both the functions $\Psi, \Phi(\cdot)$ such that $\xi \in \text{Dom}(V^{-1})$ and moreover

$$\mathbf{E}\Psi(\| V^{-1}[\xi] \|_X) < \infty, \quad (3.16)$$

or equally such that the r.v. $\| V^{-1}\xi \|_X$ belongs to the Orlicz space $L(\Psi)$.

Proof is at the same as before, except the passing from the equality (3.9) to (3.10):

$$\lim_{n \rightarrow \infty} \mathbf{E}\Phi(\| \xi - F_n[\xi] \|_c) = 0$$

denotes

$$\lim_{n \rightarrow \infty} \| \xi - F_n[\xi] \|_c \|_{L(\Psi)} = 0.$$

4 Applications to the CLT in Banach spaces.

Let again η be a (Borelian) r.v. with values in the separable Banach space X equipped with norm $\| \cdot \|_X = \| \cdot \|$. Let $\eta_i, i = 1, 2, \dots$ be independent copies of η , may be defined on some another sufficiently rich probability space. Denote

$$S(n) = n^{-1/2} \sum_{i=1}^n \eta_i. \quad (4.1)$$

By definition, the r.v. η is said to satisfy Central Limit Theorem (CLT) in the space X , write $\eta \in CLT$, if the sequence $S(n)$ converges weakly (in distribution) as $n \rightarrow \infty$ to some Gaussian distributed in this space r.v. $S(\infty)$.

Evidently, η , $S(\infty)$ are mean zero, have finite weak second moment, have identical covariation operator. Moreover, r.v. η is *pregaussian*. This imply by definition that $\mathbf{P}(S(\infty) \in X) = 1$.

There are many works about CLT in different Banach spaces, see, e.g. [8] - [17] and reference therein. The majority of aforementioned works are devoted to the finding of necessary or sufficient conditions for these Theorem.

M.Ledoux and M.Talagrand in the book [10], p.294-296, Lemma 10.1 proved that if the r.v. η satisfies CLT in the space X , then

$$\lim_{t \rightarrow \infty} t^2 \mathbf{P}(\|\eta\| > t) = 0.$$

As a consequence: for every values $p \in (0, 2) \Rightarrow \mathbf{E}\|\eta\|^p < \infty$.

Moreover, if the r.v. η satisfies CLT in the space X , then

$$\lim_{t \rightarrow \infty} t^2 \sup_n \mathbf{P}(\|S(n)\| > t) = 0,$$

$$p \in (0, 2) \Rightarrow \sup_n \mathbf{E}\|S(n)\|^p < \infty.$$

M.Ledoux and M.Talagrand [10], p.296-302 introduced and investigated also the following norm for the r.v. η :

$$\|\eta\|_{CLT(X)} \stackrel{def}{=} \sup_n \|S(n)\|_X, \quad (4.2)$$

and proved in particular that $\eta \in CLT(X) \Rightarrow \|\eta\|_{CLT} < \infty$, and that the functional $\eta \rightarrow \|\eta\|_{CLT}$ is actually (complete) norm on the linear space of all r.v. satisfying CLT in the space X .

On the other words, the finiteness of the norm $\|\eta\|_{CLT(X)}$ is *necessary* condition for the conclusion $\eta \in CLT(X)$, but it is known that this condition is'nt *sufficient*.

Proposition 4.1. Let the centered r.v. η with values in the space X is pregaussian. Suppose in addition there exists a compact linear operator $U : X \rightarrow X$ such that

$$\sup_n \|U^{-1}S(n)\| = K(\text{Law}(\eta)) < \infty. \quad (4.3)$$

Then $\eta \in CLT(X)$.

Proof. The convergence of finite-dimensional (cylindrical) distributions of $S(n)$ to ones for $S(\infty)$ is obvious. Further, we have using Tchebychev's-Markov inequality introducing the pre-compact set

$$Z(t) = U\{B_X(t)\}, \quad B_X(t) := \{x, x \in X, \|x\|_X \leq t\} :$$

$$\mathbf{P}(S(n) \notin Z(t)) \leq \frac{K(\text{Law}(\eta))}{t} \leq \epsilon \quad (4.4)$$

for sufficiently greatest values t . Thus, the sequence of distributions of the variables $\text{Law}(S(n))$ is tight in this space.

This completes the proof of proposition 4.1.

The examples of applying of this approach, i.e. by means of assertion (4.1), may be found in the articles [11]-[14].

5 Concluding remarks. Case of linear topological spaces. Open questions.

Perhaps, the main result of this report remains true for the linear topological spaces X if it is countable normed or metrizable. In contradiction, in the preprint [7] is constructed an example of linear topological space L and a Borelian r.v. ζ with values in this space such that its distribution does not has a compact embedded linear support of the full measure.

References

- [1] BULDYGIN V.V. (1984). *Supports of probabilistic measures in separable Banach spaces*. Theory Probab. Appl. **29**, v.3, pp. 528-532.
- [2] CIESIELSKI Z. *Properties of the orthogonal Franklin system*. Studia Math., **23** : **2**, (1963), pp. 141-157.
- [3] FRANKLIN P. *A set of continuous orthogonal functions*. Math. Ann., **100**, (1928), pp. 522-529.
- [4] S. KACZMARZ, H. STEINHAUS. *Theorie der Orthogonalreihen*, Chelsea, reprint (1951).
- [5] KANTOROVICZ L.V., AKILOV G.P. *Functional Analysis*. Moskow, Nauka, 1984, Third edition (in Russian).
- [6] OSTROVSKY E.I. (1980). *On the support of probabilistic measures in separable Banach spaces*. Soviet Mathematic, Doklady, v.255, No 6, pp. 836-838, (in Russian).
- [7] OSTROVSKY E. *Support of Borelian measures in separable Banach spaces*. arXiv:0808.3248v1 [math.FA] 24 Aug 2008
- [8] DUDLEY R.M. *Uniform Central Limit Theorem*. Cambridge University Press, 1999.
- [9] HEINKEL B. *Measures majorantes et le theoreme de la limite centrale dans $C(S)$* . Z. Wahrscheinlichkeitstheory. verw. Geb., (1977). **38**, 339-351.

- [10] LEDOUX M., TALAGRAND M. (1991) *Probability in Banach Spaces*. Springer, Berlin, MR 1102015.
- [11] E.OSTROVSKY, L.SIROTA. *Central Limit Theorem in Hölder spaces in the terms of majorizing measures*. arXiv:1409.6054v1 [math.PR] 21 Sep 2014
- [12] E.OSTROVSKY, L.SIROTA. *CLT for continuous random processes under approximations terms*. arXiv:1304.0250v1 [math.PR] 31 Mar 2013
- [13] E.OSTROVSKY, L.SIROTA. *Central Limit Theorem and exponential tail estimations in hybrid Lebesgue-continuous spaces*. arXiv:1309.2344v1 [math.PR] 9 Sep 2013
- [14] E.OSTROVSKY, L.SIROTA. *Central Limit Theorem and exponential tail estimations in mixed (anisotropic) Lebesgue spaces*. arXiv:1308.5606v1 [math.PR] 26 Aug 2013
- [15] RATCHKAUSKAS A, SUQUET CH. *Central limit theorems in Hölder topologies for Banach space valued random fields*. Teor. Veroyatnost. i Primenen., 2004, Volume 49, Issue 1, Pages 109-125, (in Russian).
- [16] A. RATCHKAUSKAS, CH. SUQUET. *Necessary and sufficient condition for the Hölderian functional central limit theorem*. J. Theoret. Probab. **17**, (2004), 221-243.
- [17] RATCHKAUSKAS A, SUQUET CH. *Hölder norm test statistics for epidemic change*. J. Statist. Plann. Inference, **126**, (2004), 495-520.
- [18] M.A.KRASNOSELSKY, YA.B.RUTISKY. (1961). *Convex functions and Orlicz Spaces*. P. Noordhoff LTD, The Netherland, Groningen.
- [19] M.M. RAO, Z.D.REN. (1991). *Theory of Orlicz Spaces*. Basel-New York, Marcel Decker.
- [20] M.M. RAO, Z.D.REN. (2002). *Application of Orlicz Spaces*. Basel-New York, Marcel Decker.